

***Chionea austriaca* in caves and artificial galleries of Slovenia (Diptera, Limoniidae)**

Tone NOVAK¹, Ignac SIVEC², Franc JANŽEKOVIČ¹ & Erhard CHRISTIAN³

¹ Department of Biology, Faculty of Natural Sciences and Mathematics, University of Maribor, Koroška 160, SI-2000 Maribor.

E-mail: tone.novak@uni-mb.si; franc.janzekovic@uni-mb.si

² Slovenian Museum of Natural History, Prešernova 20, SI-1000 Ljubljana.

E-mail: isivec@pms-lj.si

³ Institute of Zoology, Department of Integrative Biology and Biodiversity Research, University of Natural Resources and Applied Life Sciences, Vienna (BOKU), Gregor Mendel Str. 33, A-1180 Wien. E-mail: erhard.christian@boku.ac.at

***Chionea austriaca* in caves and artificial galleries of Slovenia (Diptera, Limoniidae).** - We sampled 54 caves and galleries in north and east Slovenia methodically for ecological data and the invertebrate fauna. The snow fly *Chionea austriaca*, recorded in 37% of the cavities, proved indifferent with regard to most of the abiotic factors, but occurred in significantly higher individual numbers in caves on north and east slopes. The subterranean sojourn of *Ch. austriaca* appears merely weather-mediated. No indications of larval development in the caves were found.

Keywords: abiotic factors - speleobiology - distribution - snow fly.

INTRODUCTION

As their scientific and trivial names imply, snow flies – species of the genus *Chionea* – are usually met on the snow surface where they can easily be noticed. Nevertheless, these wing-reduced and somewhat spiderlike limoniids spend most of their adult lifetime in leaf litter, under logs and stones, in subnivean habitats or in mammal burrows. The larvae are soil-dwelling. Low catchability makes snow flies appear so rare that Byers (1983), for instance, referred to a catch of 29 specimens as a rich yield.

Snow flies have been reported from altitudes up to 2700 m (3500 m in North America: Byers, 1983). *Chionea* (*Sphaeconophilus*) *belgica* (Becker, 1912), and *Ch. (S.) lutescens* Lundström, 1907, occur also in the colline zone and in lowlands (Grootaert, 1984; Reusch, 1988, 1997; Weber & Weidemann, 1993), whereas the bulk of the *Chionea* species prefer forest sites, or open habitats from the submontane zone upwards (Focarile, 1975).

Chionea austriaca (Christian, 1980), originally described from a Lower Austrian cave, is distributed over western Hungary (Christian, unpubl.) and parts of

Slovenia and Austria. At the northwestern and western limits of distribution *Ch. austriaca* touches the areas of two closely related species, namely *Ch. belgica* and the widespread *Ch. lutescens* (Christian, in press).

Ecological notes on European *Chionea* species focus on aboveground habitats (e.g. Heim de Balsac, 1934; Nadig, 1943, 1949; Bitsch, 1955; Sömme & Östbye, 1969; Hågvar, 1971; Mendl *et al.*, 1977; Itämies & Lindgren, 1985; Heijerman, 1987). Several authors have casually mentioned the sojourn of snow flies in caves (Strobl, 1900; Peyerimhoff, 1906; Bezzi, 1911; Czerny, 1930; Español Coll, 1955, Strinati & Aellen, 1967; Turquin, 1973; Bourne, 1977, 1979; Christian, 1980; Byers, 1983; Novak & Kuštor, 1983; Novak, 2005). Information about environmental conditions in cavities inhabited by snow flies, however, is anecdotal at the best.

The present paper addresses the abiotic conditions at sampling sites of *Ch. austriaca* in caves and artificial galleries in Slovenia. We want to test our hypothesis that this species is only loosely aligned to macrocaverns.

MATERIALS AND METHODS

During the period 1977–2001, 54 caves and artificial galleries (in the following: caves) in central and northern Slovenia (mapped in Novak, 2005) were ecologically investigated in January, April, July and October according to a standardized scheme. All accessible surfaces of the cave passages like walls, ceiling, speleothems, microhabitats below stones and wood, and water surfaces, were carefully inspected for terrestrial animals. Pitfall traps were set, on average, every 3 m starting at the entrance, and exposed for 47 ± 1 hours. Each trap was baited with 3 g of decomposing meat attached on a wire above the trapping liquid (2 cm of fruit juice with cherry and maraschino essences, and a lacing of detergent). Substrate from the close vicinity of the trap sites was sampled for Tullgren extraction.

Environmental parameters were determined chiefly according to Stewart *et al.* (1974) and Rowell (1997). The distance from a trap to entrance and surface, respectively, and the cross-section of the passage at the trap site were taken from cave maps (Novak, 2005). Illumination was measured using a Lunasix luxmeter. Substrate pH was determined with a combined electrode (91–02 Orion, USA; Orion Ionalyser 407 A) in elutriated fresh samples after 2–4 hrs stirring in an equal volume of distilled water. Substrate carbonate and organic matter contents were established for the dried and pulverized <2 mm fraction; carbonate was quantified by Scheibler calcimetry, ignition loss after heating at 700°C for 1 h. Ground and air temperatures were measured with decimal thermometers (Intos, Croatia). Air humidity was determined with a psychrometer according to August, ground moisture by drying at 105°C to constant weight. Airflow was detected with a hand-held anemometer (Munro IM159) or, when below $0.3 \text{ m}\cdot\text{s}^{-1}$, derived from the velocity of fog or magnesium smoke (cf. Andrieux, 1970).

Cold hardiness of one male, collected on a south-facing meadow slope at Studence (46°18'53" N, 15°09'51" E, 330 m) on snow, 31.01.2006, was determined in a thermostat chamber THK/V1-2020 (Elelstro, Slovenia). The specimen was enclosed in a glass tube with a scrap of wet paper and cooled down to lethal temperature in 12 hour treats in steps of 1°C/day, with intermediate 12 hour breaks at acclimatization temperature of +8°C.

Voucher specimens of *Ch. austriaca* are deposited in the Slovenian Museum of Natural History (Ljubljana), in the Muséum d'histoire naturelle de la Ville de Genève, and in the Natural History Museum, Vienna.

RESULTS

Three of 14 European *Chionea* species (Oosterbroek, 2005) have been recorded in Slovenia (Oosterbroek & Simova-Tošić, 2003, and own data; Fig. 1). *Ch. (Chionea) araneoides* Dalman, 1816 was found in the Krma valley (UTM: VM13); *Ch. (Sphaeconophilus) alpina* Bezzi, 1908 is known from Mt. Grmada (VM61, 640 m),




FIG. 1

Records of snow flies in Slovenia: ▲ *Chionea araneoides*, ○ *Ch. alpina* and ● *Ch. austriaca*.

Zdenska jama cave (VL77, 460 m), Dobropolje (VL77, 450 m), Mačkovica cave (VL47, 480 m) (Simova-Tošić & Sivec, 1977), and the alpine ice cave Ivačičeva jama in Mt. Kredarica at 2435 m a.s.l. (Novak & Kuštor, 1983); the third species, *Ch. (S.) austriaca* was encountered in numerous caves in central and eastern Slovenia (Table 1). Here we add three snow-surface records of *Ch. austriaca*: Mt. Jesenik (WM13, 688 m), Studence (WM13, 330 m) and Mt. Snežnik (VL54, 1690 m, currently the southernmost locality), and a further cave record: Brezno pod Durcami in Mt. Raduha (VM84, 1840 m, currently the highest record in Slovenia), which was not included in the systematic investigations.

In 20 (37%) of the 54 caves, 38 specimens (11 males, 27 females; sex ratio 29:71) of *Ch. austriaca* were met, at altitudes between 270 and 1515 m (Tab. 1). Tab. 2 compiles values of abiotic factors at effective trapping sites. All specimens were trapped in pitfalls exposed in January. Larvae were neither seen, nor trapped or Tullgren-extracted. Most adults occurred in the entrance zone of the caves during the winter (Fig. 2). Caves opening in northern and eastern slopes were significantly more

TABLE 1: *Chionea austriaca* in caves of Slovenia. No = eurent number as in Novak, 2005; Cad = cadastral number in the Slovene cave register; Capital letters indicate artificial galleries; UTM code (see Fig. 1); Alt = altitude (m); Bed = bedrock; e = conglomerate, l = limestone, d = dolomite, m = marble, p = pegmatite, s = schists, t = tonalite); Ind. Date = number of individuals and date of collection [ddmmyy].

Cave or gallery	No	Cad	UTM	Co-ordinates (N, E)	Alt	Bed	Ind, Date
Rački pekel	10	465	VM73	46°23'27", 14°43'14"	590	l	1 ♀, 060179
Zijalka v Dovji Gnči	11	376	VM72	46°18'18", 14°40'14"	1515	l	1 ♂, 4 ♀, 100179
Jama pri Votli peči	18	3263	VM95	46°32'51", 14°58'33"	400	p	1 ♀, 120179
Skobirjeva votlica	19	3956	VM94	46°27'45", 14°58'46"	940	d	1 ♀, 050179
Brdajsava jama	23	3497	VM90	46°07'03", 14°54'02"	651	l	1 ♂, 3 ♀, 210179
Objekt pri Žnoderu	25	B1	WM05	46°30'34", 15°03'22"	550	l	1 ♀, 110178
Rdečka jama	26	3488	WM04	46°26'54", 15°03'44"	858	e	1 ♂, 120178
Lokoviška jama	27	3959	WM03	46°22'03", 15°01'27"	370	l	1 ♀, 070178
Umetni rov v Bistriškem grabnu	30	U2	WM16	46°38'47", 15°07'56"	480	s	2 ♀, 150178
Zapečke peči	31	3208	WM15	46°32'49", 15°13'26"	610	l	1 ♂, 150178
Špegličeva jama	34	3512	WM12	46°17'58", 15°11'55"	400	l	1 ♀, 090179
Oevirkova jama v Štadlerjevem gozdu	35	5348	WM11	46°12'29", 15°11'11"	320	l	2 ♂, 2 ♀, 090179
Jama pod Herkovimi pečmi	37	1849	WM26	46°37'35", 15°16'04"	545	l	2 ♀, 190178
Jaklova luknja	38	4636	WM24	46°28'56", 15°16'49"	1100	t	1 ♂, 110178
Fantovska luknja 2	40	3967	WM21	46°12'14", 15°18'35"	480	l	1 ♂, 4 ♀, 180179
Jama v kamnolomu pri Suhem	42	4632	WM31	46°08'40", 15°23'55"	500	l	1 ♂, 18011978
Glija jama	43	84	WM30	46°06'39", 15°26'32"	515	l	1 ♀, 18011978
Luknja pri Naeeku na Planici	45	2407	WM44	46°28'30", 15°34'13"	730	m	1 ♂, 200178
Jama v kamnolomu nad Studenicami	46	252	WM42	46°17'51", 15°37'11"	400	l	1 ♂, 2 ♀, 170179
Rov	47	U4	WM41	46°09'31", 15°35'43"	270	s	1 ♀, 170179




FIG. 2

Yield of *Ch. austriaca* individuals relating to the distance between cave entrance and trap locality.

frequently settled by *Ch. austriaca* as compared to other slopes ($\chi^2 = 24.54$; $P < 0.001$), and hosted significantly more individuals ($\chi^2 = 17.05$; $P = 0.017$) (Tab. 3).

Preference for any kind of bedrock was not observed. In two cases (26 and 38 in Tab. 1) entrances were situated in various types of forests.

By comparing the meteorological conditions in caves with vs. those without *Ch. austriaca* we found that in "*Chionea* caves" the air and ground temperatures were significantly lower, and airflow significantly higher. There were no significant differences regarding relative air humidity and substrate moisture (Tab. 4). Males and females did not differ significantly with respect to any abiotic parameter at the trapping sites ($F_{1,37} = 0.000-2.614$; $P = 0.11-0.98$).

In the cold hardness test the single male of *Ch. austriaca* died at -7°C .

DISCUSSION

Adult *Ch. austriaca* were trapped in caves only during the cold season, in agreement with the period of their occurrence on snow. The temperature demands of *Ch. austriaca* are comparable to those of most other European snow flies (Wojtusiak, 1950; Bourne, 1979), and its supercooling ability – admittedly determined for one specimen only – seems to agree with the values reported for *Ch. araneoides* (mean supercooling point -7.5°C , range -10.5 / -4.5°C ; Sömme & Östbye, 1969). A vast ecological range of *Ch. austriaca* is indicated by the large altitudinal range, indifference to bedrock, its occurrence in forested as well as in non-forested areas, and inside caves by tolerance

TABLE 2: January values of abiotic factors at effective trapping sites of *Ch. austriaca*, compared to the extreme values in "*Chionea* caves" and to the mean and extreme values of all the 54 caves (based on 605-617 sampling sites).

	Mean $\pm$ s.d. (min-max) at effective trapping sites	Min-Max in " <i>Chionea</i> caves"	Mean $\pm$ s.d. (min-max) in all the 54 caves
Distance from entrance [m]	8 $\pm$ 10 (0-49)	0-121	31 $\pm$ 31 (0-121)
Distance from surface [m]	9 $\pm$ 8 (1-40)	0-70	18 $\pm$ 14 (0-80)
Passage cross-section [m ²]	5 $\pm$ 7 (0.736)	0.2-80	13 $\pm$ 31 (0.2-460)
Illumination [lx]	258 $\pm$ 169 (0~10,000)	0~100,000	194 $\pm$ 171 (0~100,000)
pH	7.9 $\pm$ 0.8 (4.3-8.6)	4.3-8.7	8.1 $\pm$ 0.6 (2.4-8.9)
CO ₃ ⁻² [%]	21.4 $\pm$ 24.0 (0.0-76.0)	0-96.6	31.9 $\pm$ 29.8 (0.0-98.7)
Ignition loss [%]	13.2 $\pm$ 15.2 (0.3-52.9)	0.3-85.9	10.7 $\pm$ 13.2 (0.3-85.9)
Air temperature [°C]	1.2 $\pm$ 3.8 (-5.0-8.9)	-5.0-11.5	3.6 $\pm$ 4.4 (-6.0-11.5)
Ground temperature [°C]	1.6 $\pm$ 3.3 (-3.0-9.0)	-4.6-12.6	4.1 $\pm$ 4.2 (-7.2-12.6)
Relative air humidity [%]	91 $\pm$ 12 (65-100)	54-100	95 $\pm$ 9 (54-100)
Substrate moisture [%]	18.4 $\pm$ 12.0 (0.2-46.7)	0.0-73.6	20.8 $\pm$ 14.7 (0.0-87.6)
Airflow [m/s]	0.07 $\pm$ 0.04 (0.00-2.00)	0.00-10.00	0.11 $\pm$ 0.73 (0.00-10.00)
Soil colour / modus (min-max)	10YR 5/4 (5Y 3/2-7.5YR 5/6) 5YR 5/8-5Y 5/2 10YR4/4 (5YR 3/2-2.5Y 6/4)		

TABLE 3: Distribution of *Ch. austriaca* with respect to slope orientation of cave entrances.

Slope orientation	<i>Ch. austriaca</i> individuals N (%)	Caves with <i>Ch. austriaca</i> N (%)	All caves N (%)
N	12 (31.6)	4 (20.0)	10 (18.5)
NE	2 (5.3)	2 (10.0)	3 (5.6)
E	12 (31.6)	6 (30.0)	14 (25.9)
SE	5 (13.2)	3 (15.0)	4 (7.4)
S	4 (10.5)	4 (20.0)	7 (13.0)
SW	0 (0.0)	0 (0.0)	4 (7.4)
W	0 (0.0)	0 (0.0)	10 (18.5)
NW	3 (7.9)	1 (5.0)	2 (3.7)
total	38 (100)	20 (100)	54 (100)

TABLE 4: Selected meteorological parameters of caves with *Ch. austriaca* present vs. absent. January values from all trap sites along the first 10 m (i.e., those without *Chionea* yield included). Significant differences asterisked *.

	<i>Ch. austriaca</i> present (N = 103)	absent (N = 159)	F _{1, 260}	P
Air temperature	4.8 $\pm$ 2.4 (0.6-9.5)	6.2 $\pm$ 2.0 (0.3-11.4)	24.59	<0.001*
Ground temperature	4.7 $\pm$ 2.5 (-0.3-11.3)	5.7 $\pm$ 2.0 (0.5-11.0)	12.78	<0.001*
Air humidity	94 $\pm$ 8 (69-100)	92 $\pm$ 22 (43-100)	2.52	0.113
Substrate moisture	23.3 $\pm$ 16.1 0.0-65.4	24.8 $\pm$ 17.0 (1.1-79.2)	0.52	0.473
Airflow	0.08 $\pm$ 0.05 (0.0-10.00)	0.06 $\pm$ 0.04 (0.0-5.00)	4.97	0.026*

of wide ranges of illumination, substrate pH, carbonate contents and ignition loss. The conspicuous prevalence of individuals in caves on north and east facing slopes contrasts with the impression that snow flies in open-air habitats do not display any slope orientation preference. Their surface activity, however, is strongly weather-

dependent inasmuch heavy frost and wind drive them into more moderate hideaways. So the disproportionately high individual numbers of *Ch. austriaca* in caves opening out on north and east slopes is not a consequence of psychrophily, but rather the result of more frequent chill avoidance migrations there. The highest trapping yield of snow flies in the entrance zone of the caves and in passages of small cross-section supports this conclusion. Lower ground temperature and higher airflow in “*Chionea* caves” may be merely the consequence of the harsh climate conditions on north and east slopes.

Wing reduction of snow flies has been explained as a result of regressive evolution due to temporary exposure to cold and windy weather (Hackman, 1964; Byers, 1961, 1983). This morphological character could lead to consider snow flies as being pre-disposed to life in subterranean habitats (cf. Vandel, 1964; Ginot & Decou, 1982). However, *Ch. austriaca* – like several other European congeners – inhabits macro-caverns facultatively and only temporarily during its epigeal ecophase in the cold season. The fact that maximum density in caves is observed in January, when virtually all females are ready for egg-laying (Christian unpubl.), does not necessarily indicate subterranean oviposition. On the one hand there is no report on larval development of *Chionea* in caves, and on the other hand an extremely long oviposition period (from November to March) was shown for *Ch. araneoides* (Hågvar, 1976), so that females of *Ch. austriaca* – let’s assume an adult life span of two months (cf. Byers, 1983) – may lay eggs after they have quitted the cave shelter. Thus, in terms of the classical biospeleological classification (Vandel, 1964), *Ch. austriaca* is best included in the troglomenes.

ACKNOWLEDGEMENTS

We thank Valika Kuštor, Tomaž Prošek and Mojmir Štangelj for help with the field investigations, Dušan Devetak (Dept. Biol. Univ. Maribor) and the anonymous reviewers for insightful remarks on the manuscript, and Michelle Gadpaille (Dept. Angl. Amer. Univ. Maribor) for linguistic improvements. The study was partly supported by the Slovene Ministry of High Education, Science and Technology within the Biodiversity research programme (P1-0078).

REFERENCES

- ALLEN, S. E., GRIMSHAW, H. M., PARKINSON, J. A. & QUARMBY, C. 1974. Chemical analysis of ecological materials. *Blackwell Scientific Publications, Oxford*, 565 pp.
- ANDRIEUX, C. 1970. Contribution à l'étude du climat des cavités naturelles des massifs karstiques. *Annales de Spéléologie* 25: 441-559.
- BEZZI, M. 1911. Diptères (première série), suivi d'un Appendice sur les Diptères cavernicoles recueillis par le Dr Absolon dans les Balkans. *Biospeologica*, 20. *Archives de Zoologie Expérimentale et Générale* 8(5): 1-87.
- BITSCH, J. 1955. Le genre *Chionea* (Diptera, Tipulidae). *Ecologie, systématique et morphologie. Travaux du Laboratoire de Zoologie et de la Station Agricole Grimaldi de la Faculté des Sciences de Dijon* 11: 1-26.
- BOURNE, J. D. 1977. Remarques sur la présence du genre *Niphadobata* (= *Chionea*) à basse altitude (Dipt. Tipulidae). *Mitteilungen der schweizerischen entomologischen Gesellschaft* 50: 338.

- BOURNE, J. D. 1979. Description de trois nouvelles espèces du genre *Niphadobata* (Diptera: Tipulidae), avec quelques remarques biogéographiques. *Revue suisse de Zoologie* 86: 233-249.
- BYERS, G. W. 1961. Biology and classification of *Chionea* (Diptera: Tipulidae). XI. *Internationaler Kongress für Entomologie, Wien 1960, Verhandlungen* 1: 188-191.
- BYERS, G. W. 1983. The crane fly genus *Chionea* in North America. *The University of Kansas Science Bulletin* 52(6): 59-195.
- CHRISTIAN, E. 1980. Eine neue Schneefliege aus der *Chionea*-Verwandtschaft: *Niphadobata austriaca* n. sp. (Dipt.: Tipulidae). *Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen* 32: 77-79.
- CHRISTIAN, E. (in press). Snow flies of the genus *Chionea* (Diptera: Limoniidae) in Austria. *Contributions to Natural History, Bern*.
- CZERNY, L. 1930. Dipteren auf Schnee und in Höhlen. *Mitteilungen des Königlichen naturwissenschaftlichen Instituts Sofia* 3: 113-118.
- ESPAÑOL COLL, F. 1955. Nuevos datos sobre la entomofauna cavernícola de la provincia de Lérida. *Eos* 31: 261-273.
- FOCARILE, A. 1975: *Chionea minuta* Tahv. specie boreo-alpina nuova per la fauna italiana e terza serie di dati sulla geonemia di *Chionea alpina* Bezzi (Diptera Limnobiidae). *Atti della Accademia delle Scienze di Torino. Classe di Scienze Fisiche, Matematiche e Naturali* 109: 135-144.
- GINET, R. & DECOU, V. 1982. Initiation à la biologie et l'écologie souterraines. *Delarge, Paris*, 345 pp.
- GROOTAERT, P. 1984. Redescription of *Niphadobata belgica* (Becker, 1912) comb. nov. (Diptera - Tipulidae), a snowfly from low altitudes in Belgium. *Annales de la Société Royale Zoologique de Belgique* 114: 241-247.
- HACKMAN, W. 1964. On reduction and loss of wings in Diptera. *Notulae Entomologicae* 44: 73-93.
- HÄGVAR, S. 1971. Field observations on the ecology of a snow insect, *Chionea araneoides* Dalm. (Dipt., Tipulidae). *Norsk Entomologisk Tidsskrift* 18: 33-37.
- HÄGVAR, S. 1976. Phenology of egg development and egg-laying in a winter-active insect, *Chionea araneoides* Dalm. (Dipt., Tipulidae). *Norwegian Journal of Entomology* 23: 193-195.
- HEIJERMAN, T. 1987. De sneeuwmug *Niphadobata belgica* (BECKER, 1912) in Nederland (Diptera: Tipulidae). *Entomologische Berichten, Amsterdam*, 47: 113-118.
- HEIM DE BALSAC, H. 1934. Un Diptère peu connu de la faune française: *Chionea lutescens* Lund. (Dipt. Tipulidae). Précisions sur sa biologie. *Bulletin de la Société entomologique de France* 39: 102-105.
- ITÄMIES, J. & LINDGREN, E. 1985. The ecology of *Chionea* species (Diptera, Tipulidae). *Notulae Entomologicae* 65: 29-31.
- MENDL, H., MÜLLER, K. & VIRAMO, J. 1977. Vorkommen und Verbreitung von *Chionea araneoides* Dalm., *C. crassipes* Boh. und *C. lutescens* Lundstr. (Diptera, Tipulidae) in Nord-europa. *Notulae Entomologicae* 57: 85-90.
- NADIG, A. 1943. Beiträge zur Kenntnis der Dipterengattung *Chionea* (I. Teil). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 19: 53-65.
- NADIG, A. 1949. Beiträge zur Kenntnis der Dipterengattung *Chionea*. III. Teil: Die Faktoren, welche für das Vorkommen auf dem Schnee massgebend sind. *Mitteilungen der schweizerischen entomologischen Gesellschaft* 22: 323-345.
- NOVAK, T. & KUŠTOR, V. 1983. On the ecology of an alpine ice cave. *Mémoires de biospéologie* 10: 117-125.
- NOVAK, T. 2005. Terrestrial fauna from cavities in Northern and Central Slovenia, and a review of systematically ecologically investigated cavities. *Acta carsologica, Ljubljana*, 34: 169-210.

- OOSTERBROEK, P. 2005. Genus *Chionea*.- Fauna Europaea, version 1.2, <http://www.faunaeur.org>.
- OOSTERBROEK, P. & SIMOVA-TOŠIĆ, D. 2003. An annotated list of the species of Pediciidae, Limoniidae and Cylindrotomidae of Slovenia, Croatia, Bosnia-Herzegovina, Serbia, Montenegro and Macedonia, with additions for the Tipulidae (Insecta, Diptera). *Studia dipterologica, Halle (Saale)*, 10: 441-452.
- PEYERIMHOFF, P. 1906. Recherches sur la faune cavernicole de Basses-Alpes. *Annales de la Société Entomologique de France* 35: 203-211.
- REUSCH, H. 1988. *Niphadobata belgica* (Becker, 1912) neu für Deutschland (Diptera: Limoniidae). *Entomologische Zeitschrift* 98: 300-304.
- REUSCH, H. 1997. Notes on *Chionea* (*Sphaeconophilus*) species in central Europe (Diptera: Limoniidae). *Bulletin de la Société Neuchâteloise des Sciences Naturelles* 120: 169-173.
- ROWELL, D. L. 1997. Bodenkunde. Untersuchungsmethoden und ihre Anwendungen. *Springer, Berlin etc.*, 614 pp.
- SIMOVA-TOŠIĆ, D. & SIVEC, I. 1977. Dve retke vrste limonide (Diptera) iz Slovenije. *Acta entomologica Jugoslavica* 13: 45-49.
- SÖMME, L. & ÖSTBYE, E. 1969. Cold-hardiness in some winter active insects. *Norsk Entomologisk Tidsskrift* 16: 45-48.
- STRINATI, P. & AELLEN, V. 1967. Faune actuelle de deux grottes suisses de haute altitude. *Rassegna Speleologica Italiana* 19: 84-87.
- STROBL, G. 1900. Dipterenfauna von Bosnien, Hercegovina und Dalmatien. *Wissenschaftliche Mittheilungen aus Bosnien und der Hercegowina, Sarajevo*, 7: 552-670.
- TURQUIN, M.-J. 1973. La colonisation de quelques grottes du Jura par *Niphadobata alpina* Bezzi (Dipt. Tipulidae). *International Journal of Speleology* 5: 21-28.
- VANDEL, A. 1964. Biospéologie. *Gauthier-Villars, Paris*, 619 pp.
- WEBER, G. & WEIDEMANN, G. 1993. *Niphadobata lutescens* (Lundström, 1908) and *N. belgica* (Becker, 1912) from the Solling Mountains, Lower Saxony, FRG (Diptera, Limoniidae). *Deutsche Entomologische Zeitschrift* 40: 259-263.
- WOJTUSIAK, H. 1950. The temperature preferendum of winter insects of the genus *Boreus* (Panorpatae) and *Chionea* (Diptera). *Bulletin international, Polska Akademia Umiejętnosci, Wydział matematyczno-przyrodniczy, Série B*, 11: 125-143.